arXiv:1411.6281v1 [physics.plasm-ph] 23 Nov 2014

Comment on the paper 'An analytic functional form for characterization and generation of axisymmetric plasma boundaries', PPCF 55 (2013) 095009

Ap Kuiroukidis[1] and G. N. Throumoulopoulos[2]

[1]Technological Education Institute of Serres 621 24 Serres, Greece

[2]Physics Department, University of Ioannina, GR 451 10 Ioannina, Greece

E-mail: kouirouki@astro.auth.gr, gthroum@uoi.gr

In Ref. [1] it was proposed an analytic form to describe the boundary of an axisymmetric plasma. The form determines analytically smooth boundaries, e.g. D-shaped ones, and numerically boundaries having an X-point. Here we propose an alternative form describing boundaries with X-points fully analytically.

The two coordinates are (R, z) and they are normalized with respect to the major radius R_0. For example for ITER we have $R_0 = 6.2m$. The boundary is up-down asymmetric and it consists of a smooth upper part and a lower part possessing an X-point. For the upper part we have four parameters. One is the inverse aspect ratio $\epsilon_0 = a/R_0$. For example in the case of ITER it is $a = 2.1m$ and thus $\epsilon_0 = 0.338$. The other three are the upper elongation κ, the upper triangularity δ and a parameter n related to the steepness of the triangularity of the upper boundary curve, i.e. to the mean value of the derivative $d\rho/d\theta$ [see Eqs. (1) and (2) below]. For the ITER case it is $\kappa = 1.86$ and $\delta = 0.5$. Introducing the normalized coordinates $\rho = R/R_0$ and $\zeta = z/R_0$, we have for the ζ coordinate of the uppermost point (see Fig. 1): $\zeta_u = \kappa\epsilon_0$ and $\delta = (1 - \rho_\delta)/\epsilon_0$.

The form for the upper part of the boundary is given by

$$\begin{aligned} \rho &= 1 + \epsilon_0 cos(\tau + \alpha sin(\tau)) \\ \zeta &= \zeta_u sin(\tau) \end{aligned} \tag{1}$$

where $\alpha = sin^{-1}(\delta)$. Thus the following relations hold: $\rho_\delta = 1 - \delta\epsilon_0$ and $\theta_\delta = \pi - tan^{-1}(\kappa/\delta)$. The parameter τ is any increasing function of the usual polar angle θ, satisfying $\tau(0) = 0$, $\tau(\pi) = \pi$ and $\tau(\theta_\delta) = \pi/2$. In our form we have taken

$$\begin{aligned} \tau(\theta) &= t_0\theta^2 + t_1\theta^n \\ t_0 &= \frac{\theta_\delta^n - \frac{1}{2}\pi^n}{\pi\theta_\delta^n - \theta_\delta^2\pi^{n-1}} \\ t_1 &= \frac{-\theta_\delta^2 + \frac{1}{2}\pi^2}{\pi\theta_\delta^n - \theta_\delta^2\pi^{n-1}} \end{aligned} \tag{2}$$

The uppermost point of the bounding curve has coordinates (ρ_δ, ζ_u).



Figure 1. The boundary of an axisymmetric plasma having a saddle point, P_X, as determined by Eqs. (1), (3) and (4) containing six shaping parameters.

Now we consider the lower part of the boundary which is located at the $\zeta < 0$-part of the plane. It depends on two additional parameters, namely the lower elongation $\bar{\kappa}$ and the lower triangularity $\bar{\delta}$. In terms of these parameters we have $\zeta_d = \bar{\kappa}\epsilon_0$ and $\bar{\theta}_\delta = \pi - tan^{-1}(\bar{\kappa}/\bar{\delta})$, where $-\zeta_d$ corresponds to the X-point. The equation for the left-lower part of the bounding curve, with respect to the X-point, is given by

$$\begin{aligned} \rho_l &= 1 + \epsilon_0 cos(\theta) \\ \zeta_l &= -[2p_1\epsilon_0(1+cos(\theta)]^{1/2} \\ p_1 &= \frac{\zeta_d^2}{2\epsilon_0(1+cos\bar{\theta}_\delta)}, \quad (\pi \le \theta \le 2\pi - \bar{\theta}_\delta) \end{aligned} \tag{3}$$

while that for the right-lower part is given by

$$\begin{aligned} \rho_r &= 1 + \epsilon_0 cos(\theta) \\ \zeta_r &= -[2p_2\epsilon_0(1-cos(\theta)]^{1/2} \\ p_2 &= \frac{\zeta_d^2}{2\epsilon_0(1-cos\bar{\theta}_\delta)}, \quad (2\pi - \bar{\theta}_\delta \le \theta \le 2\pi) \end{aligned} \tag{4}$$

The coordinates of the X-point are $(\rho_X, \zeta_X) = (1+\epsilon_0 cos\bar{\theta}_\delta, -\zeta_d)$. At the midplane $\zeta = 0$ the upper and the right- and left-lower parts of the boundary meet smoothly, that is the derivative $d\zeta/d\rho$ is continuous there.

In summary, alternatively to the numerical functional-form determination of the boundary of an axisymmetric plasma having an X-point proposed in Ref. [1], we propose

here a completely analytic form for boundaries of this kind. This form contains six free parameters associated with the boundary shaping and, as the case of Ref. [1], should be useful for studies of fusion plasmas.

Aknowledgments

This work was supported by (a) the National Programme for the Controlled Thermonuclear Fusion, Hellenic Republic, (b) the European Union's Horizon 2020 research and innovation programme under grant agreement number 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

References

[1] T C Luce, Plasma Phys. Control. Fusion **55** (2013) 095009; idid **55** (2013) 119501 (corrigendum).